

Distance-balanced closure of some graphs

N. Ghareghani ^{b,d,1,2}, B. Manoochehrian ^c, M. Mohammad-Noori ^{a,d}

^a*Department of Mathematics, Statistics and Computer Science, University of Tehran*

P.O. Box 14155-6455, Tehran, Iran

^b*Department of Mathematical Sciences, K.N. Toosi University of Technology*

^c*Academic Center for Education, Culture and Research (ACECR), Tehran Branch, Tehran, Iran*

^d*School of Mathematics, Institute for Research in Fundamental Sciences (IPM),*

P.O. Box 19395-5746, Tehran, Iran

Emails: ghareghani@ipm.ir, behzad@khayam.ut.ac.ir,

morteza@ipm.ir, mnoori@khayam.ut.ac.ir

December 20, 2010

Abstract

In this paper we prove that any distance-balanced graph G with $\Delta(G) \geq |V(G)| - 3$ is regular. Also we define notion of distance-balanced closure of a graph and we find distance-balanced closures of trees T with $\Delta(T) \geq |V(T)| - 3$.

AMS Classification: 05B20; 05E30

Keywords: Distances in graphs; distance-balanced graphs; distance-balanced closure

1 Introduction

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. We denote $|V(G)|$ by n . The set of neighbors of a vertex $v \in V(G)$ is denoted by $N_G(v)$, and $N_G[v] = N_G(v) \cup \{v\}$. The degree of a vertex v is denoted by $\deg_G(v)$ and minimum degree and maximum degree of G denoted by $\delta(G)$ and $\Delta(G)$, respectively. The distance $d_G(u, v)$ between vertices u and v is the length of a shortest path between u and v in G . The diameter $\text{diam}(G)$ of graph G is defined as $\max\{d_G(u, v) : u, v \in V(G)\}$.

¹Corresponding author

²The research of the first author was in part supported by a grant from IPM (No. 89050046)

For an edge xy of a graph G W_{xy}^G is the set of vertices closer to x than y , more formally

$$W_{xy}^G = \{u \in V(G) | d_G(u, x) < d_G(u, y)\}.$$

Moreover, ${}_xW_y^G$ is the set of vertices of G that have equal distances to x and y or

$${}_xW_y^G = \{u \in V(G) | d_G(u, x) = d_G(u, y)\}.$$

These sets play important roles in metric graph theory, see for instance [1, 2, 3, 4]. The set W_{xy}^G also appears in chemical graph theory to compute the well-investigated Szeged index of a graph (see [5, 6]), defined as:

$$Sz(G) = \sum_{xy \in E(G)} |W_{xy}^G| \cdot |W_{yx}^G|.$$

Distance-balanced graphs are introduced in [7] as graphs for which $|W_{xy}^G| = |W_{yx}^G|$ for every pair of adjacent vertices $x, y \in V(G)$.

In [7], the parameter $b(G)$ of a graph G is introduced as the smallest number of the edges which can be added to G such that the obtained graph is distance-balanced. Since the complete graph is distance-balanced, this parameter is well-defined. We call graph G a *distance-balanced closure* of H if H is a spanning subgraph of G and $|E(G)| = b(H) + |E(H)|$; in other words, a distance-balanced closure of H is a graph G which contains H as a spanning subgraph and has minimum number of edges. As mentioned in [7], the computation of $b(G)$ is quite hard in general but it might be interesting in some special cases. In this paper we compute $b(G)$ for all trees T with $\Delta(T) \geq |V(T)| - 3$. In Section 2, we compute that distance-balanced closure of graphs G with $\Delta(G) = n - 1$. In Section 3, and Section 4, we concern graphs G with $\Delta(G) = n - 2$ and $\Delta(G) = n - 3$, respectively. Then we compute $b(T)$ for all trees T with $\Delta(T) = n - 2$ and $\Delta(T) = n - 3$.

Here we mention some more definitions and notations about trees. Let P_n denoted the path with n vertices. A tree which has exactly one vertex of degree greater than two is said to be *starlike*. The vertex of maximum degree is called the *central vertex*. We denote by $S(n_1, n_2, \dots, n_k)$ a starlike tree in which removing the central vertex leaves disjoint paths $P_{n_1}, P_{n_2}, \dots, P_{n_k}$. We say that $S(n_1, n_2, \dots, n_k)$ has branches of length $n_1, n_2, \dots, n_k$. It is obvious that $S(n_1, n_2, \dots, n_k)$ has $n_1 + n_2 + \dots + n_k + 1$ vertices. For simplicity a starlike with α_i branches of length n_i ($1 \leq i \leq k$) is denoted by $S(n_1^{\alpha_1}, n_2^{\alpha_2}, \dots, n_k^{\alpha_k})$.

2 Distance-balanced graphs with maximum degree $n - 1$

In this section we prove for any graph G with $\Delta(G) = n - 1$, the only distance-balanced closure of G is the complete graph K_n . The following result is very useful in this paper. It is in fact a slight modification of Corollary 2.3 of [7].

Theorem 1. *Let G be a graph with diameter at most 2 and H be a distance-balanced graph such that G is a spanning subgraph of H . Then H is a regular graph. Moreover, every regular graph with diameter at most 2 is distance-balanced.*

Corollary 2. *For every integer $n \geq 1$, the graph $K_{1,m}$ has a unique distance-balanced closure which is isomorphic to K_{m+1} , hence, $b(K_{1,m}) = \binom{m+1}{2} - m$.*

Proof. Let G be a distance-balanced closure of $K_{1,m}$. By Theorem 1, G is a regular graph and since $K_{1,m}$ has a vertex of degree m , G should be m -regular, hence $G \cong K_{m+1}$. $\square$

The following is an immediate conclusion of Theorem 1.

Corollary 3. *Let G be a graph with n vertices and $\Delta(G) = n - 1$. Then the graph G has a unique closure which is isomorphic to K_n .*

3 Distance-balanced graphs with maximum degree $n - 2$

In this section, we prove that any distance-balanced graph G with $\Delta(G) = n - 2$ is a regular graph, using this we construct a distance-balanced closure of G where G is a tree with this property and then compute $b(G)$.

The following Lemma will be used occasionally in this paper and the proof is easily deduced from the definition of W_{xy}^G .

Lemma 4. *Let x and y be two vertices of a graph G , then $W_{xy}^G \cap N_G(y) = \emptyset$. Furthermore, $N_G(y) \setminus W_{yx}^G \subseteq N_G(x)$.*

Theorem 5. *Let $G = S(2, 1^{m-1})$ be a starlike tree and H be a distance-balanced graph containing G as a spanning subgraph. Then $\text{diam}(H) \leq 2$, hence, H is an r -regular graph for some $m \leq r \leq m + 1$.*

Proof. Suppose that the vertices of G are labeled as shown in Figure 1.




Figure 1

If $oy \in E(H)$, then H contains $K_{1,m+1}$ as a spanning subgraph, so by Theorem 1 and Corollary 2, $H \cong K_{m+2}$.

So, we may assume that $oy \notin E(H)$ (and consequently $\text{diam}(H) \neq 1$). We prove $\text{diam}(H) = 2$; For this, it is enough to show that $d(y, x_i) \leq 2$ for $i = 1, \dots, m$. Let i be an integer with $1 \leq i \leq m$; From $|W_{x_i o}^H| = |W_{o x_i}^H|$ and Lemma 4, we conclude that $W_{x_i o}^H \subseteq \{y\}$; We may suppose $W_{x_i o}^H = \emptyset$ (Otherwise, $W_{x_i o}^H = \{y\}$ leads to $x_i y \in E(H)$ and $d_H(x_i, y) = 1$). By Lemma 4, this leads to

$$N_H(x_i) = \{o, x_1, x_2, \dots, x_m\} \setminus \{x_i\},$$

from which, we conclude that $x_1 \in N_H(x_i) \cap N_H(y)$ and $d_H(x_i, y) = 2$. Hence, $\text{diam}(H) = 2$, as required. The result is now concluded from Theorem 1. $\square$

Theorem 6. *Let $G = S(2, 1^{m-1})$ be a starlike of order $m + 2$. Then*

$$b(G) = \begin{cases} \frac{m^2}{2} - 1 & \text{if } m \text{ is even;} \\ \binom{m+1}{2} & \text{otherwise.} \end{cases}$$

Proof. Let the vertices of G be labeled as in Figure 1, and $\overline{G}$ be a distance-balanced closure of G . First, suppose that m is an odd integer; Since there is no m -regular graph of order $m + 2$, by Theorem 5, $\overline{G} \cong K_{m+2}$.

Now, suppose that m is an even integer. Let $H = K_{m+2}$ be a complete graph with vertex set $V(G)$ and M be a complete matching of H which contains the edge oy . Then $H \setminus M$, is an m -regular graph with diameter 2. Hence, by Theorem 1 and Theorem 5, $\overline{G} = H \setminus M$, is a distance-balanced closure of G and $b(G) = \frac{m^2}{2} - 1$. $\square$

Corollary 7. *Let G be a connected graph of order n , with $\Delta(G) = n - 2$ and H be a distance-balanced graph which contains G as a spanning subgraph. Then H is either an $(n - 2)$ -regular graph or the complete graph K_n .*

Proof. In this case $S(2, 1^{n-3})$ is an spanning subgraph of G . So, by Theorem 5, H is either an $(n - 2)$ -regular graph or the complete graph K_n . $\square$

4 Distance-balanced graphs with maximum degree $n - 3$

In this section we will prove that every distance-balanced graph with $\Delta(G) = n - 3$ is regular. Moreover, by constructing distance-balanced closure of trees with $\Delta(T) = n - 3$ we compute $b(T)$ for these trees.

Theorem 8. Let $G = S(2^2, 1^{m-2})$ be a starlike of order $m+3$ and H be a distance-balanced graph which contains G as a spanning subgraph. Then $\text{diam}(H) \leq 2$, hence, H is an r -regular graph with $r \geq m$.

Proof. Suppose the vertices of G are labeled as in Figure 2.




Figure 2

If either oy or oz be an edge of H , then by Theorem 5, H is an r -regular graph with $r \geq m+1$, which proves this theorem. So, suppose that $oy, oz \notin E(H)$.

By Lemma 4, for each $1 \leq i \leq m$, $W_{x_i o}^H \subseteq \{y, z\}$. Now, we prove that $\deg_H(x_i) = m$, for each $i = 1, \dots, m$; For this, we consider three possible cases:

Case 1. $|W_{x_i o}^H| = 0$. Then $W_{x_i o}^H = \emptyset$ and $W_{o x_i}^H = \emptyset$. Hence, by Lemma 4, $N_H(x_i) = \{o, x_1, x_2, \dots, x_m\} \setminus \{x_i\}$ and $\deg_H(x_i) = m$.

Case 2. $|W_{x_i o}^H| = 1$. Without loss of generality we can assume that $W_{x_i o}^H = \{y\}$. Then there is an integer $1 \leq j \leq m$ such that $W_{o x_i}^H = \{x_j\}$. Since $d_H(o, y) = 2$, $yx_i \in E(H)$. Hence, using Lemma 4, $N_H(x_i) = \{o, y, x_1, x_2, \dots, x_m\} \setminus \{x_j\}$ and $\deg_H(x_i) = m$.

Case 3. $|W_{x_i o}^H| = 2$. We have $W_{x_i o}^H = \{y, z\}$. Since $d_H(o, y) = d_H(o, z) = 2$, we conclude $yx_i, zx_i \in E(H)$. Since $|W_{o x_i}^H| = 2$, there are integers j and k such that $W_{o x_i}^H = \{x_j, x_k\}$.

Hence, by Lemma 4, we have $N_H(x_i) = \{o, y, z, x_1, x_2, x_3, \dots, x_m\} \setminus \{x_i, x_j, x_k\}$ and $\deg_H(x_i) = m$.

Next, we prove that $\deg_H(y) \geq m-3$ and $\deg_H(z) \geq m-3$. From $\deg_H(x_1) = m$ and Lemma 4, it concludes that $|W_{y x_1}^H| \leq 2$, hence, $|W_{x_1 y}^H| \leq 2$, which means that there are at most two elements in $N_H(x_1) \setminus N_H[y]$. Using this and Lemma 4, we provide $\deg_H(y) \geq m-3$. With a similar argument, the inequality $\deg_H(z) \geq m-3$ is concluded.

Now, by using $\deg_H(y), \deg_H(z) \geq m-3$, $\deg_H(x_i) \geq m$, ($i = 1, \dots, m$), and $oy, oz \notin E(H)$, we conclude that every two nonadjacent vertices have a common neighbor, provided that $m \geq 7$. This means that $\text{diam}(H) = 2$, which proves the result in case $m \geq 7$, using Theorem 1. For cases

$3 \leq m \leq 6$, through a case by case inspection (by using $\deg_H(x_i) \geq m$, $i = 1, \dots, m$), the same result is obtained. $\square$

Theorem 9. For the starlike tree $G = S(2^2, 1^{m-2})$ of order $m + 3$, $b(G) = \frac{m^2+m-4}{2}$.

Proof. Let the vertices of G be labeled as in Figure 2 and $\overline{G}$ be a distance-balanced closure of G . Now, we are going to construct $\overline{G}$. Let $H = K_{m+3}$ be a complete graph with the same vertex set as H . Omit the edges of cycles $C_1 = x_1x_2x_3 \dots x_mx_1$ and $C_2 = oyzo$ from H to obtain $\overline{G} = H \setminus (C_1 \cup C_2)$. Now, $\overline{G}$ is an m -regular graph with diameter 2, which contains G as a spanning subgraph, so by Theorem 8 and Theorem 1, $\overline{G}$ is a distance-balanced closure of G and $b(G) = \frac{m^2+m-4}{2}$. $\square$

Theorem 10. Let G be the graph of Figure 3 and H be a distance-balanced graph which contains G as a spanning subgraph. Then $\text{diam}(H) \leq 2$, hence, H is a regular graph. Moreover, $b(G) = \frac{m^2+m-4}{2}$.




Figure 3

Proof. If either $oy \in E(H)$ or $oz \in E(H)$, then by Theorem 5, $\text{diam}(H) \leq 2$ and H is a regular graph. So, suppose that neither oy nor oz is in $E(H)$. Since $|W_{x_iy}^H| = |W_{yx_i}^H|$ and $o \in W_{x_iy}^H$, there exists a vertex x_i , $i \neq 1$, such that $yx_i \in E(H)$. Therefore, graph H contains graph $S(2^2, 1^{m-2})$ as a spanning subgraph and using Theorem 8, $\text{diam}(H) \leq 2$ and H is a regular graph. Furthermore, the graph introduced in the proof of Theorem 9, is also distance-balanced closure of G . Hence $b(G) = \frac{m^2+m-4}{2}$. $\square$

Theorem 11. Consider the starlike tree $G = S(3, 1^{m-1})$ of order $m + 3$ and let H be a distance-balanced graph which contains G as a spanning subgraph. Then H is an r -regular graph for some $m \leq r \leq m + 2$.

Proof. Let the vertices of G be labeled as in Figure 4.




Figure 4

If either $oy \in E(H)$ or $oz \in E(H)$, then by Theorem 5, $\text{diam}(H) \leq 2$ and H is a regular graph. Suppose that $oy \notin E(H)$ and $oz \notin E(H)$. Since $|W_{yz}^H| = |W_{zy}^H|$, the vertex z is adjacent with at least one vertex in $\{x_1, x_2, \dots, x_m\}$. Hence, H contains either the graph of Figure 3 or the graph $S(2^2, 1^{m-2})$, as a spanning subgraph. So, by Theorem 8 and Theorem 10, $\text{diam}H \leq 2$ and H is a regular graph, as desired. $\square$

Corollary 12. *Let G be a connected graph of order n with $\Delta(G) = n - 3$. Then every distance-balanced graph H which contains G as a spanning subgraph, is regular.*

Proof. Since $\Delta(G) = n - 3$, G contains at least one of the graphs $S(2^2, 1^{n-2})$, $S(3, 1^{n-1})$ or the graph shown in Figure 3, as a spanning subgraph. Hence, the result follows from Theorem 9, Theorem 13 and Theorem 10. $\square$

Theorem 13. *For the starlike tree $G = S(3, 1^{m-1})$ of order $m + 3$, $b(G) = \frac{m^2+m-4}{2}$.*

Proof. Let the vertices of G be labeled as in Figure 4 and let $\overline{G}$ be a distance-balanced closure of G . Now, we are going to construct $\overline{G}$. Let $H = K_{m+3}$ be a complete graph with the same vertex set as G . Omit the edges of cycles $C_1 = x_3x_4 \dots x_mx_3$ and $C_2 = oxyx_2x_1zo$ from H to obtain $\overline{G} = H \setminus (C_1 \cup C_2)$. Then the graph $\overline{G}$ is an n -regular graph with diameter 2, which contains G as a spanning subgraph. So by Theorem 11, $\overline{G}$ is a distance-balanced closure of G and $b(G) = \frac{m^2+m-4}{2}$. $\square$

Conclusion. In previous sections, we have proved that any connected distance-balanced graph G with $\Delta(G) \geq |V(G)| - 3$ is a regular graph, moreover, distanced-closure of such a graph G is a smallest regular graph which contains G . This helped us to find a distance-balanced closure of trees T with $\Delta(T) \geq |V(T)| - 3$ and to compute $b(T)$ for such trees.

References

- [1] H. J. Bandelt and V. Chepoi, Metric graph theory and geometry: a survey, manuscript, 2004.
- [2] V. Chepoi, Isometric subgraphs of hamming graphs and d -convexity, *Cybernetics* **24** (1988), 6-10 (Russian, English transl.)
- [3] D. ž. Dijoković, Dictance preserving subgraphs of hypercubes, *J. Combin. Theory Ser B.* **14** (1973), 263-267.
- [4] D. Eppstein, The lattice dimation of a graph, *European J. Combin.* **26** (2005), 585-592.
- [5] A. Graovac, M. Juvan, M.Petkovšek, A. Vasel and J. Žerovnik, The Szged index of fascia-graphs, *MATCH Commun. Math. Comput. Chem.* **49** (2003), 47-66.
- [6] I. Gutman, L. Popovič, P. V. Khadikar S. Karmarkar, S. Joshi and M. Mandloi, Relations between Wiener and Szged indices of monocyclic molecules, *MATCH Commun. Math. Comput. Chem.* **35** (1997), 91-103.
- [7] J. Jerebic, S. Klavžar, D. F. Rall, Dictance-balanced graphs, *Ann. Combin.* **12** (2008), 71-79.